

IDEALS WHICH GENERALIZE (v^0)

PIOTR KALEMBA AND SZYMON PLEWIK

ABSTRACT. We consider ideals $d^0(\mathcal{V})$ which are generalizations of the ideal (v^0) . We formulate counterparts of Hadamard's theorem. Then, adopting the base tree theorem and applying Kulpa-Szymański Theorem, we obtain $cov(d^0(\mathcal{V})) \leq add(d^0(\mathcal{V}))^+$.

1. INTRODUCTION

Let ω denotes the set of all natural numbers and $|X|$ denotes the cardinality of a set X . If $\mathcal{F}$ is a family of sets, then $add(\mathcal{F})$ is the least cardinality of a subfamily $\mathcal{G} \subseteq \mathcal{F}$ such that the union $\bigcup \mathcal{G}$ is not in $\mathcal{F}$, but $cov(\mathcal{F})$ is the least cardinality of a subfamily $\mathcal{G} \subseteq \mathcal{F}$ such that $\bigcup \mathcal{G} = \bigcup \mathcal{F}$. Also, $d^0(\mathcal{F})$ denotes an ideal defined as follows. A set $X \in d^0(\mathcal{F})$, whenever $X \subset \bigcup \mathcal{F}$ and for each $V \in \mathcal{F}$ there exists $U \in \mathcal{F}$ such that $U \subseteq V$ and $U \cap X = \emptyset$.

In this note, products $\prod X_i$ are examined where $i \in \omega$. In fact, it is assumed that each X_i is a finite discrete space with more than one point. For infinite X_i , we make some comments, only. Considerations are related to some trees which we call trimmed trees. Each trimmed tree $T[A, \alpha]$ is uniquely determined by two parameters $A \in [\omega]^\omega$ and $\alpha \in \prod X_i$, and it is a subset of the union $\bigcup \{X_0 \times X_1 \times \dots \times X_n : n \in \omega\} = \prod_{fin} X_i$. As usual, elements of a trimmed tree $T[A, \alpha] \subseteq \prod_{fin} X_i$ are called nodes. Unions of nodes which belong to $\prod X_i$ are called branches. The family of all branches of a trimmed tree is the perfect subset $[T[A, \alpha]] \subseteq \prod X_i$. So, our terminology is standard, compare [5] or [7].

2000 *Mathematics Subject Classification.* Primary: 03E05; Secondary: 03E35, 06A06.

Key words and phrases. Base tree; Fusion relation; Trimmed tree; $add(d^0(\mathcal{V}))$; $cov(d^0(\mathcal{V}))$.

We are guided by the notion $J_f(\alpha)$ which was considered by G. Moran and D. Strauss [10]. For a given function f defined on $\prod X_i$ and a fixed point $\alpha \in \prod X_i$ the family $J_f(\alpha) \subseteq [\omega]^\omega$ consists of all sets $A \in [\omega]^\omega$ such that f is constant on the set of all functions $\beta \in \prod X_i$ such that $\alpha(n) = \beta(n)$ for $n \in \omega \setminus A$. There are conditions sufficient for some $J_f(\alpha)$ to be non-empty, see [10] Theorems 1 - 3. In fact, $A \in J_f(\alpha)$ says that the function f is constant on the perfect set $[T[A, \alpha]]$.

In [3], [6] or [9] were considered properties of the ideal (v^0) . Ideals $d^0(\{[T] : T \in \mathcal{V}\})$ are generalizations of (v^0) . In fact, (v^0) is $d^0(\{[T] : T \in \mathcal{V}\})$, where it is assumed that $X_n = \{0, 1\}$ for all $n \in \omega$. Here, we generalize properties of the ideal (v^0) for sets X_n being finite and with more than one point. Albeit, we do not consider properties which depend on the assumption that sets X_n have less than k -points for a fixed $k \in \omega$, compare Theorem 3 in [10], or that the set $\{n : |X_n| = k\}$ is finite for each $k \in \omega$, compare Example 1 in [10]. We adopt the base tree theorem, see [1] and [2]. Applying so called Kulpa-Szymański Theorem, we obtain the final result of this paper, i.e. inequalities $add(d^0(\mathcal{V})) \leq cov(d^0(\mathcal{V})) \leq add(d^0(\mathcal{V}))^+$ under the hypothesis that factors X_i of a product $\prod X_i$ are finite and have more than one point.

2. TRIMMED TREES

Suppose $X_0, X_1, \dots$ be an infinite sequence of sets with more than one point. Let $\prod X_i$ denotes the Cartesian product of theses sets. Fix a function $\alpha \in \prod X_i$ and an infinite subset $A \subseteq \omega$. The subset

$$\{\beta \in \prod X_i : \alpha|_{\omega \setminus A} = \beta|_{\omega \setminus A}\}$$

is a perfect subset of $\prod X_i$ equipped with product topology, where each X_n is considered with the discrete topology. We will denote this subset $[T[A, \alpha]]$, since it can be described as the set of all infinite branches of the tree which we call a *trimmed tree*.

Any trimmed tree $T[A, \alpha] \subseteq \prod_{fin} X_i$ is the union of sets of nodes $T[A, \alpha, n]$, where $n \in \omega$. We assume that the empty set is a node of any $T[A, \alpha]$, too. One obtains sets of nodes $T[A, \alpha, n]$ by the following procedure $T[A, \alpha, n] = \{s \cup \{(n, \alpha(n))\} : s \in T[A, \alpha, n-1]\}$, whenever $n \notin A$, but $T[A, \alpha, n] = \{s \cup \{(n, x)\} : s \in T[A, \alpha, n-1] \text{ and } x \in X_n\}$ for $n \in A$. This procedure is inductive, so we assume that $T[A, \alpha, 0] = \{(0, \alpha(0))\}$, whenever $0 \notin A$, and $T[A, \alpha, 0] = \{(0, x) : x \in X_0\}$,

for $0 \in A$. In consequence $T[A, \alpha] = \bigcup \{T[A, \alpha, n] : n \in \omega\} \cup \{\emptyset\}$ and $[T[A, \alpha]] = \{\beta \in \prod X_i : \alpha|_{\omega \setminus A} = \beta|_{\omega \setminus A}\}$.

Obviously, $T[A, \alpha]$ does not depend on the restriction $\alpha|_A$. Also, $T[A, \alpha] = T[B, \beta]$ implies $\alpha|_{\omega \setminus A} = \beta|_{\omega \setminus B}$, what means $A = B$ and $\alpha(n) = \beta(n)$ for $n \in \omega \setminus A$.

Suppose $\prod X_i$ is fixed. Put

$$\mathcal{V} = \{T[A, \alpha] : A \in [\omega]^\omega \text{ and } \alpha \in \prod X_i\}.$$

If $\alpha \in \prod X_i$ and $s \in \prod_{fin} X_i$, then let α_s denotes the unique function $\beta \in \prod X_i$ such that $s \subset \beta \subseteq \alpha \cup s$. If $Y \subseteq \prod X_i$ and $s \in \prod_{fin} X_i$, then Y_s denotes the set $\{\alpha_s : \alpha \in Y\}$, but Y^* denotes the family of all functions α_s , where $\alpha \in Y$ and $s \in \prod_{fin} X_i$. Thus

$$Y^* = \bigcup \{Y_s : s \in \prod_{fin} X_i\}.$$

Finally, put $\mathcal{V}^* = \{[T]^* : T \in \mathcal{V}\}$. From now on, we assume that a Cartesian product $\prod X_i$ is fixed where each X_i is a set with more than one point. This will be applied to symbols $\mathcal{V}$ or $\mathcal{V}^*$, always. A few facts will need that each set X_i is finite, additionally.

Lemma 1. *Each member of $\mathcal{V}^*$ contains continuum many pairwise disjoint members of $\mathcal{V}^*$.*

Proof. Fix $[T]^* \in \mathcal{V}^*$, where $T = T[A, \alpha]$. Let $\mathcal{R}(A)$ be an almost disjoint family of the cardinality continuum consisting of infinite subsets of A . Fix a function $\gamma \in \prod X_i$ such that $\gamma(n) \neq \alpha(n)$, for any $n \in A$. For each $C \in \mathcal{R}(A)$ choose an infinite subset $V_C \subset C$ such that $C \setminus V_C$ is infinite, too. Let $\alpha_C \in \prod X_i$ be a function such that

$$\alpha_C(n) = \begin{cases} \gamma(n), & \text{for } n \in C; \\ \alpha(n), & \text{for } n \notin C. \end{cases}$$

The family $\{[T[V_C, \alpha_C]]^* : C \in \mathcal{R}(A)\}$ is a desired one. $\square$

If $T \in \mathcal{V}$ and $s \in \prod_{fin} X_i$, then T_s denotes the tree $\{\alpha_s|n : \alpha \in [T] \text{ and } n \in \omega\}$. Note that, notions T_s and Y_s are used in different contexts. Each tree T_s consists of nodes, but any Y_s consists of infinite sequences. We have assumed that each X_n has at least two points, hence any tree $T[A, \alpha]$ has continuum many branches. If $T = T[A, \alpha]$

and $s \in \prod_{fin} X_i$, then $T_s = T[A \setminus |s|, \alpha_s]$. Therefore, $T \in \mathcal{V}$ implies $T_s \in \mathcal{V}$.

3. FUSION RELATIONS

Let $a_0, a_1, \dots$ and $b_0, b_1, \dots$ be increasing enumerations of all points of $A = \{a_n : n \in \omega\} \in [\omega]^\omega$ and $B \in [\omega]^\omega$, respectively. Put

$$T[A, \alpha] \subseteq_n T[B, \beta],$$

whenever $T[A, \alpha] \subseteq T[B, \beta]$ and $a_0 = b_0, a_1 = b_1, \dots, a_n = b_n$. Thus, the decreasing sequence of relations $(\subseteq_n)_{n \in \omega}$ is defined. These relations hold between elements of $\mathcal{V}$. Always, $\subseteq_{n+1}$ is contained in $\subseteq_n$. So, we can apply the method of fusion, compare [5], using these relation to trimmed trees. In many papers, facts about fusion are presented without proof. Since details considered here are not so obvious, we run full proofs here.

Lemma 2. *Let $(T[A_n, \alpha_n])_{n \in \omega}$ be a sequence of elements of $\mathcal{V}$. If always $T[A_{n+1}, \alpha_{n+1}] \subseteq_n T[A_n, \alpha_n]$, then there exists $C \in [\omega]^\omega$ and $\alpha \in \prod X_i$ such that $T[C, \alpha] \subseteq T[A_n, \alpha_n]$ for any $n \in \omega$.*

Proof. An inclusion $T[A_{n+1}, \alpha_{n+1}] \subseteq T[A_n, \alpha_n]$ implies $A_{n+1} \subseteq A_n$ and $\alpha_n|_{\omega \setminus A_n} = \alpha_{n+1}|_{\omega \setminus A_n}$ and $\alpha_n|_{\omega \setminus A_n} = \alpha_{n+1}|_{\omega \setminus A_n} \subseteq \alpha_{n+1}|_{\omega \setminus A_{n+1}}$. Hence, the union $\bigcup \{\alpha_n|_{\omega \setminus A_n} : n \in \omega\}$ is a function. Fix $\alpha \in \prod X_i$ which extends this union. Functions α and α_n coincide on the set $\omega \setminus A_n$, thus $T[A_n, \alpha] = T[A_n, \alpha_n]$. Let $a_0^n, a_1^n, \dots$ be the increasing enumeration of points of A_n . Put $C = \{a_n^n : n \in \omega\}$. If $k \geq n$, then $a_k^k \in A_k \subseteq A_n$. If $k < n$, then

$$T[A_n, \alpha_n] = T[A_n, \alpha] \subseteq_{n-1} T[A_{n-1}, \alpha] \subseteq_{n-2} \dots \subseteq_k T[A_k, \alpha],$$

hence $T[A_n, \alpha] \subseteq_k T[A_k, \alpha]$. This implies $a_k^k \in A_n$. Therefore $C \subseteq A_n$. Finally, $T[C, \alpha] \subseteq T[A_n, \alpha] = T[A_n, \alpha_n]$ for any $n \in \omega$. $\square$

From now on, the ideal $d^0(\{[T] : T \in \mathcal{V}\})$ will be shortly denoted $d^0(\mathcal{V})$.

Lemma 3. *Let $s, t \in \prod_{fin} X_i$ and $Y \subseteq \prod X_i$ and assume that $|X_k| < \text{add}(d^0(\mathcal{V}))$ for any $k \in \omega$. Then, $Y_s \in d^0(\mathcal{V})$ if and only, if $Y_t \in d^0(\mathcal{V})$.*

Proof. Suppose $|s| = |t|$. Fix $T \in \mathcal{V}$. Let $Y_s \in d^0(\mathcal{V})$. If $t \notin T$, then $[T] \cap Y_t = \emptyset$. If $t \in T$, then choose $P_s \in \mathcal{V}$ such that $[P_s] \cap Y_s = \emptyset$ and $P_s \subseteq T_s$. Hence, $P_t \subseteq T_t \subseteq T$ and $[P_t] \cap Y_t = \emptyset$. Therefore $Y_t \in d^0(\mathcal{V})$.

Suppose $s \in \prod_{fin} X_i$ and $Y_s \in d^0(\mathcal{V})$. Since Y is a subset of the union $\bigcup \{Y_u : u \in \prod_{fin} X_i \text{ and } |u| = |s|\}$, then it is contained in an union of less than $add(d^0(\mathcal{V}))$ many elements of $d^0(\mathcal{V})$. Hence $Y \in d^0(\mathcal{V})$.

Now, suppose that $Y \in d^0(\mathcal{V})$ and $s \in \prod_{fin} X_i$. If $t \in \prod_{fin} X_i$ and $|s| = |t|$, then $\{\alpha \in Y : \alpha|_{|s|} = t\} = Y_t \cap Y = (Y_t \cap Y)_t \subseteq Y$. Thus $(Y_t \cap Y)_s \in d^0(\mathcal{V})$. Also, $Y_s = \bigcup \{(Y_t \cap Y)_s : t \in \prod_{fin} X_i \text{ and } |s| = |t|\}$. Hence Y_s is an union of less than $add(d^0(\mathcal{V}))$ many elements of $d^0(\mathcal{V})$. Therefore $Y_s \in d^0(\mathcal{V})$. $\square$

Lemma 4. *Suppose that $|X_i| < add(d^0(\mathcal{V}))$ for all $i \in \omega$. Let $k \in \omega$ and $T \in \mathcal{V}$. If $Y \in d^0(\mathcal{V})$, then there exists a tree $P \in \mathcal{V}$ such that $P \subseteq_k T$ and $[P] \cap Y = \emptyset$.*

Proof. Let $T = T[A, \alpha]$ and $a_0, a_1, \dots$ be the increasing enumeration of all points of A . Consider the union $U = \bigcup \{Y_s : s \in X_0 \times \dots \times X_{a_k}\}$. It consists of less than $add(d^0(\mathcal{V}))$ many sets, each one from $d^0(\mathcal{V})$ by Lemma 3. Thus $U \in d^0(\mathcal{V})$. Take $Q \in \mathcal{V}$ such that $Q \subseteq T$ and $[Q] \cap U = \emptyset$. Check that $[Q_s] \cap Y = \emptyset$ for any $s \in X_0 \times \dots \times X_{a_k}$. Finally put $P = \bigcup \{Q_s : s \in T \cap X_0 \times \dots \times X_{a_k}\}$. $\square$

We do not know, whether lemmas 2, 3 and 4 are valid for $\prod X_i = \omega^\omega$. Their proofs would work if $d^0(\mathcal{V})$ was a σ -ideal. If all sets X_i are finite, then hypotheses of these lemmas are fulfilled. Moreover, then $add(d^0(\mathcal{V}))$ is an uncountable cardinal.

Theorem 5. *If each X_i is a finite set, then $d^0(\mathcal{V})$ is a σ -ideal.*

Proof. Assume $S_0, S_1, \dots$ is an increasing sequence of elements of the ideal $d^0(\mathcal{V})$. Fix $T \in \mathcal{V}$ and put $T_0 = T$. Using Lemma 4, choose inductively trees $T_k \in \mathcal{V}$ such that $T_{k+1} \subseteq_k T_k$ and $[T_k] \cap S_k = \emptyset$. Thus, it has been defined a sequence of elements of $\mathcal{V}$ satisfying hypotheses of Lemma 2. So, there exists a tree $P \in \mathcal{V}$ such that $P \subseteq T$ and $[P] \cap \bigcup_{k \in \omega} S_k = \emptyset$. $\square$

Corollary 6. *If each X_i is a finite set and $Y \in d^0(\mathcal{V})$, then $Y^* \in d^0(\mathcal{V})$.*

Proof. By Lemma 3, $Y_s \in d^0(\mathcal{V})$ for each $s \in \prod_{fin} X_i$. Since $Y^* = \bigcup \{Y_s : s \in \prod_{fin} X_i\}$, it is a countable union of elements of $d^0(\mathcal{V})$. Thus $Y^* \in d^0(\mathcal{V})$ by Theorem 5. $\square$

4. COUNTERPARTS OF HADAMARD'S THEOREM

Two sequences of countable sets $(a_n)_{n \in \omega}$ and $(b_n)_{n \in \omega}$ form a (ω, ω) -gap, whenever

$$a_0 \subset^* a_1 \subset^* \dots a_n \subset^* \dots \subset^* b_n \subset^* \dots \subset^* b_1 \subset^* b_0$$

and no set c fulfills $a_n \subset^* c \subset^* b_n$ for all $n \in \omega$. The famous Hadamard's theorem says that there are no (ω, ω) -gaps, compare [4] or [11]. This theorem can be formulated in our's notations. Indeed, assume that always $X_i = \{0, 1\}$ and identify each subset $Y \subseteq \omega$ with its characteristic function which belongs to $\prod X_i$. Then one can check that Hadamard's theorem is equivalent to the property that any decreasing sequence of elements of $\mathcal{V}^*$ has a lower bound. Theorem 8 extends this property.

If $T = T[A, \alpha] \in \mathcal{V}$, then the tree T is determined by the function

$$\delta(T[A, \alpha])(n) = \begin{cases} \{\alpha(n)\}, & \text{whenever } n \notin A; \\ X_n, & \text{whenever } n \in A. \end{cases}$$

For a tree $P \in \mathcal{V}$, we have $\beta \in [P]$ if and only, if $\beta(n) \in \delta(P)(n)$ for each $n \in \omega$. Also, $\beta \in [P]^*$ if and only, if $\beta(n) \in \delta(P)(n)$ for all, but finitely many $n \in \omega$. One can check that, if $[P]^*, [T]^* \in \mathcal{V}^*$, then $[P]^* \subseteq [T]^*$ if and only, if $\delta(P)(k) \subseteq \delta(T)(k)$ for all, but finitely many $k \in \omega$.

Lemma 7. *If $P, T \in \mathcal{V}$ and $[P]^* \subseteq [T]^*$, then for each $n \in \omega$ there exists a tree $Q \in \mathcal{V}$ such that $Q \subseteq_n T$ and $[Q]^* = [P]^*$.*

Proof. Fix $k_0 \in \omega$ such that $\delta(P)(k) \subseteq \delta(T)(k)$, for any $k \geq k_0$. If $T = T[A, \alpha]$, then let a_n be the n -th element of A . Put

$$\delta(Q)(m) = \begin{cases} \delta(T)(m), & \text{for } m \leq \max\{a_n, k_0\}, \\ \delta(P)(m), & \text{for } m > \max\{a_n, k_0\}. \end{cases}$$

The function $\delta(Q)$ uniquely determines the tree $Q \in \mathcal{V}$ which is a desired one. $\square$

Theorem 8. *Let $(W_n)_{n \in \omega}$ be a sequence of elements of $\mathcal{V}^*$. If $W_{n+1} \subseteq W_n$ for any $n \in \omega$, then there exists $W \in \mathcal{V}^*$ such that $W \subseteq W_n$, for any $n \in \omega$.*

Proof. Choose $T_n \in \mathcal{V}$ such that $W_n = [T_n]^*$. Inductively, construct a sequence of trees $(Q_n)_{n \in \omega}$ such that $Q_{n+1} \subseteq_n Q_n \in \mathcal{V}$ and $[Q_n]^* = [T_n]^*$, using Lemma 7. By Lemma 2, there exists $T \in \mathcal{V}$ such that $T \subseteq Q_n$ and $W = [T]^* \subseteq [Q_n]^* = [T_n]^* = W_n$, for all $n \in \omega$. $\square$

Note that, Lemma 7 and Theorem 8 do not require that sets X_i are finite. Theorem 8 immediately follows that $d^0(\mathcal{V}^*)$ is a σ -ideal.

Corollary 9. *If each X_i is a finite set, then $d^0(\mathcal{V}) = d^0(\mathcal{V}^*)$.*

Proof. If $Y \in d^0(\mathcal{V})$ and $[T]^* \in \mathcal{V}^*$, then $Y^* \in d^0(\mathcal{V})$ by Corollary 6. There exists a tree $P \subseteq T$ such that $[P] \cap Y^* = \emptyset$. Hence $[P]^* \cap Y^* = \emptyset$ and $[P]^* \cap Y = \emptyset$ and finally $Y \in d^0(\mathcal{V}^*)$.

If $Y \in d^0(\mathcal{V}^*)$ and $T \in \mathcal{V}$, then there exists $[P]^* \subseteq [T]^*$ such that $[P]^* \cap Y = \emptyset$. Therefore $[P] \cap Y = \emptyset$. By Lemma 7 one can assume that $P \subseteq T$, so $Y \in d^0(\mathcal{V})$. $\square$

Families $\mathcal{V}$ and $\mathcal{V}^*$ are not isomorphic with respect to the inclusion. Any decreasing sequence of elements of $\mathcal{V}^*$ has a lower bound, see Theorem 8. But a sequence of trees $(T[\omega \setminus n, \alpha])_{n \in \omega}$ has no lower bound in $\mathcal{V}$.

5. A VERSION OF BASE TREE

Base Matrix Lemma, see [1], or Base Matrix Tree, compare [2] or [6], are adopted to trimmed trees in this part. We omit some proofs, since they are completely analogical to these which are in [1], [2], [6], etc. From now on, assume that all X_i are countable. Elements $[Q]^*, [P]^* \in \mathcal{V}^*$ are *incompatible* whenever the intersection $[P]^* \cap [Q]^*$ contains no element of $\mathcal{V}^*$. Thus, if $[Q]^*$ and $[P]^*$ are incompatible, then $[P]^* \cap [Q]^*$ is countable. But, if $[P]^* \cap [Q]^*$ is uncountable, then $[P]^* \cap [T]^* \in \mathcal{V}^*$. Indeed, the intersection $[P]^* \cap [Q]^*$ is a countable union of closed sets $[P_s] \cap [Q_t]$, where $s, t \in \prod_{fin} X_i$. Therefore, some $[P_s] \cap [Q_t]$ has to be

uncountable and hence $P_s \cap Q_t$ is a trimmed tree, moreover $[P_s \cap Q_t]^* = [P]^* \cap [Q]^*$.

A family $\mathcal{P}$ is called *v-partition*, whenever $\mathcal{P}$ is a maximal family, with respect to the inclusion, of pairwise incompatible elements of $\mathcal{V}^*$. Any collection of *v-partitions* is called *v-matrix*. We say that a *v-partition* $\mathcal{P}$ refines a *v-partition* $\mathcal{Q}$ (briefly $\mathcal{P} \prec \mathcal{Q}$), if for every $[P]^* \in \mathcal{P}$ there exists $[Q]^* \in \mathcal{Q}$ such that $[P]^* \subseteq [Q]^*$. A matrix $\mathcal{H}$ is called *shattering* if for any $[T]^* \in \mathcal{V}^*$ there exists a *v-partition* $\mathcal{P} \in \mathcal{H}$ such that at least two elements of $\mathcal{P}$ are compatible with $[T]^*$. The least cardinality of a shattering matrix we denote $\kappa(\prod X_i)$.

The poset $(\mathcal{V}^*, \subseteq)$ is separative, i.e. if $[P]^*$ is not contained in $[T]^*$, then there exists $[Q]^* \in \mathcal{V}^*$ such that $[Q]^* \subseteq [P]^*$ and $[Q]^*$ is incompatible with $[T]^*$. Indeed, if $[P]^* \setminus [T]^* \neq \emptyset$, then the set $Z = \{n \in \omega : \delta(P)(n) \setminus \delta(T)(n) \neq \emptyset\}$ is infinite. Fix a set $N \in [Z]^\omega$ such that there exist infinitely many $n \in \omega \setminus N$ for, which $\delta(P)(n) = X_n$. Fix also, a function α with domain N such that $\alpha(n) \in \delta(P)(n) \setminus \delta(T)(n)$ for $n \in N$. Put

$$\delta(Q)(n) = \begin{cases} \delta(P)(n), & \text{for } n \notin N; \\ \{\alpha(n)\}, & \text{for } n \in N. \end{cases}$$

The function $\delta(Q)$ uniquely determines the tree $Q \in \mathcal{V}$ which is a disered one.

For any tree $T \in \mathcal{V}$ a poset

$$(\{[P]^* \in \mathcal{V}^* : [P]^* \subseteq [T]^*\}, \subseteq)$$

is isomorphic with the poset $(\mathcal{V}^*, \subseteq)$. Moreover, posets

$$(\{P \in \mathcal{V} : P \subseteq T\}, \subseteq) \text{ and } (\mathcal{V}, \subseteq)$$

are isomorphic, too. Additionally, if $T = T[A, \alpha]$, then each bijection $\phi : A \rightarrow \omega$ determines these isomorphisms. Indeed, for $T[B, \beta] \subseteq T[A, \alpha]$, put $\Phi(T[B, \beta]) = T[\phi(B), \beta \circ \phi^{-1}]$. If $T[C, \gamma] \in \mathcal{V}$, then put $\Psi(T[C, \gamma]) = T[\phi^{-1}(C), \gamma \circ \phi \cup \alpha|_{\omega \setminus A}] \subseteq T[A, \alpha]$. We have

$$\Phi \circ \Psi(T[C, \gamma]) = T[C, \gamma] \text{ and } \Psi \circ \Phi(T[B, \beta]) = T[B, \beta].$$

Thus Φ and Ψ are mutually inverse bijections. Moreover, Φ and Ψ preserve the relation of inclusion between trees, hence they are isomorphisms. Now, we can define isomorphism Φ^* of posets

$$(\{[P]^* \in \mathcal{V}^* : [P]^* \subseteq [T]^*\}, \subseteq) \text{ and } (\mathcal{V}^*, \subseteq)$$

as follows $\Phi^*([P]^*) = [\Phi(P)]^*$, for any $[P]^* \subseteq [T]^*$ and $[P]^* \in \mathcal{V}^*$.

The next lemma is a counterpart of the lemma 2.6 in [1].

Lemma 10. *If $\mathcal{H}$ is a v -matrix of the cardinality less than $\kappa(\prod X_i)$, then there exists a v -partition $\mathcal{P}$ which refines each v -partition $\mathcal{Q} \in \mathcal{H}$.*

Proof. Orders $(\{[Q]^* \in \mathcal{V}^* : [Q]^* \subseteq [T]^*\}, \subseteq)$ and $(\mathcal{V}^*, \subseteq)$ are isomorphic and separative. So, one obtains a proof analogous to the proof of the lemma 2.6 in [1]. $\square$

Theorem 11. $\kappa(\prod X_i)$ is a regular uncountable cardinal number.

Proof. Using Theorem 8, one can proceed analogous as in [1] (corollaries 2.8 and 2.9) or as in [2] (Proposition 3.3). $\square$

Theorem 12. *There exists a v -matrix $\mathcal{H} = \{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}$ such that if $\alpha < \beta < \kappa(\prod X_i)$, then $\mathcal{P}_\beta \prec \mathcal{P}_\alpha$. Moreover, for any $[T]^* \in \mathcal{V}^*$ there exists $[P]^* \in \bigcup \mathcal{H}$ where $[P]^* \subseteq [T]^*$.*

Proof. A proof is completely analogous to the proof of Lemma 2.11 in [1], or to the proof of Theorem 3.4 in [2]. $\square$

6. CARDINAL INVARIANTS

In this part results hold for the σ -ideal $d^0(\mathcal{V}^*)$. One can check $d^0(\mathcal{V}^*) \subseteq d^0(\mathcal{V})$ similarly as in the proof of Corollary 9. We do not know whether $d^0(\mathcal{V}^*) = d^0(\mathcal{V})$. To obtain results for the ideal $d^0(\mathcal{V})$ we have to assume that sets X_i are finite.

Lemma 13. *If $\mathcal{P}$ is a v -partition, then the complement of the union $\bigcup \mathcal{P}$ belongs to $d^0(\mathcal{V}^*)$.*

Proof. If $[T]^* \in \mathcal{V}^*$, then take $[P]^* \in \mathcal{P}$ such that $[P]^* \cap [T]^* \in \mathcal{V}^*$. Since $[P]^* \cap [T]^* \subseteq \bigcup \mathcal{P}$ and $[P]^* \cap [T]^* \subseteq [T]^*$, we are done. $\square$

Lemma 14. *If $S \in d^0(\mathcal{V}^*)$, then there exists a v -partition $\mathcal{P}$ such that $\bigcup \mathcal{P} \cap S = \emptyset$.*

Proof. For each $[T]^* \in \mathcal{V}^*$ fix $[P]^* \in \mathcal{V}^*$ such that $[P]^* \subseteq [T]^*$ and $[P]^* \cap S = \emptyset$. Any v -partition consisting of just fixed $[P]^*$ is a desired one. $\square$

Observe that, if $\mathcal{P}$ is a v -partition and $S \subseteq \prod X_i$ is a selector of $\mathcal{P}$, i.e. $S \cap [P]^*$ has exactly one point for each $[P]^* \in \mathcal{P}$, then $S \in d^0(\mathcal{V})$. Indeed, for any $[T]^* \in \mathcal{V}^*$, there exist $[P]^* \in \mathcal{P}$ such that $[T]^* \cap [P]^* \in \mathcal{V}^*$. Then $[T]^* \cap [P]^* \cap S$ has no more than one point. By Lemma 1, there exists $[Q]^* \in \mathcal{V}^*$ which is disjoint with S and such that $[Q]^* \subseteq [T]^*$.

Theorem 15. $\kappa(\prod X_i) = \text{add}(d^0(\mathcal{V}^*))$.

Proof. Let $\mathcal{F} \subseteq d^0(\mathcal{V}^*)$ and $|\mathcal{F}| < \kappa(\prod X_i)$. By Lemma 14, for each $W \in \mathcal{F}$ choose a v -partition $\mathcal{P}_W$ such that $\bigcup \mathcal{P}_W \cap W = \emptyset$. By Lemma 10 there exists a v -partition $\mathcal{P}$ which refines each v -partition $\mathcal{P}_W$ for $W \in \mathcal{F}$. The set $\prod X_i \setminus \bigcup \mathcal{P}$ is element of $d^0(\mathcal{V}^*)$ and contains $\bigcup \mathcal{F}$. Thus, we have showed that $\text{add}(d^0(\mathcal{V}^*)) \geq \kappa(\prod X_i)$.

Let $\{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}$ be a v -matrix like in the Theorem 12. We will construct inductively v -matrix $\{\mathcal{Q}_\alpha : \alpha < \kappa(\prod X_i)\}$ such that for any $\alpha < \kappa(\prod X_i)$, $\mathcal{Q}_\alpha \prec \mathcal{P}_\alpha$ and if $V \in \mathcal{Q}_\alpha$, then $V \setminus \bigcup \mathcal{Q}_{\alpha+1} \neq \emptyset$. Suppose, that we have already defined a v -partition $\mathcal{Q}_\beta$ and assume that $\alpha = \beta + 1 < \kappa(\prod X_i)$. Fix a selector S of v -partition $\mathcal{Q}_\beta$. By Lemma 14 there exists a v -partition $\mathcal{Q}$ such that $\bigcup \mathcal{Q} \cap S = \emptyset$. Let $\mathcal{Q}_\alpha$ be any v -partition which refines the v -partition $\mathcal{Q}$ and the v -partition $\mathcal{P}_\alpha$. The remaining inductive steps are obvious. For any $T \in \mathcal{V}$ the intersection $[T]^* \cap \bigcup \{\prod X_i \setminus \bigcup \mathcal{Q}_\alpha : \alpha < \kappa(\prod X_i)\}$ is nonempty. Indeed, for any $T \in \mathcal{V}$ choose $\alpha < \kappa(\prod X_i)$ and $[P]^* \in \mathcal{Q}_\alpha$ such that $[P]^* \subseteq [T]^*$. Thus $\emptyset \neq [P]^* \setminus \bigcup \mathcal{Q}_{\alpha+1} \subseteq \bigcup \{\prod X_i \setminus \bigcup \mathcal{Q}_\alpha : \alpha < \kappa(\prod X_i)\}$. Therefore, the family $\{\prod X_i \setminus \bigcup \mathcal{Q}_\alpha : \alpha < \kappa(\prod X_i)\}$ witnesses that $\text{add}(d^0(\mathcal{V}^*)) \leq \kappa(\prod X_i)$. $\square$

Theorem 16. If all sets X_i are countable, then $\omega_1 \leq \kappa(\prod X_i) = \text{add}(d^0(\mathcal{V}^*)) \leq \text{cov}(d^0(\mathcal{V}^*)) \leq \text{cf}(\mathfrak{c})$.

Proof. By Lemma 1 each set $S \subseteq \prod X_i$ of the cardinality less than continuum belongs to $d^0(\mathcal{V})$. Choose sets $S_\alpha \subseteq \prod X_i$ for $\alpha < \text{cf}(\mathfrak{c})$ such that $\prod X_i = \bigcup \{S_\alpha : \alpha < \text{cf}(\mathfrak{c})\}$ and $|S_\alpha| < \mathfrak{c}$. Thus $\text{cov}(d^0(\mathcal{V})) \leq \text{cf}(\mathfrak{c})$. $\square$

If $\mathcal{P}$ is a v -partition, then one can choose subsets $N_C \subset C \in \mathcal{P}$ of the cardinality less than $\mathfrak{c}$ such that sets $A \setminus N_A$ and $B \setminus N_B$ are disjoint for any distinct members A, B of $\mathcal{P}$. One can do this by the induction using the fact that the intersection of any distinct members of $\mathcal{P}$ is countable. But, if $\{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}$ is a v -matrix like in Theorem 12, then put

$M_C = \bigcup \{N_{C_\beta} : \beta \leq \alpha\}$ whenever $C \in \mathcal{P}_\alpha$ and $C \subseteq C_\beta \in \mathcal{P}_\beta$. If all sets X_i are countable, then sets M_C have cardinalities less than the continuum, by Theorem 16. Under such assumptions, the family $\{C \setminus M_C : C \in \bigcup \{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}\}$ is called *base matrix*. Any base matrix consists of sets which are either disjoint or one is contained in the other. The topology on $\prod X_i$ generated by a base matrix is called *matrix topology*. So, the family

$$\{\prod X_i\} \cup \{C \setminus M_C : C \in \bigcup \{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}\}$$

is a base for the matrix topology.

Lemma 17. *Suppose sets X_i are countable. A subset $Y \subset \prod X_i$ is nowhere dense with respect to a matrix topology if and only, if $Y \in d^0(\mathcal{V}^*)$.*

Proof. Fix a base matrix

$$\mathcal{B} = \{C \setminus M_C : C \in \bigcup \{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}\}.$$

At first, let $Y \subset \prod X_i$ be a nowhere dense with respect to the matrix topology. Fix a set $V \in \mathcal{V}^*$ and choose $W \subseteq V$ such that $W \in \mathcal{B}$ and $W \cap Y = \emptyset$. Since $W = C \setminus M_C$, where $C \in \mathcal{V}^*$ and $|M_C| < \mathfrak{c}$, the set W contains some $D \in \mathcal{V}^*$. Any such D witnesses that $Y \in d^0(\mathcal{V}^*)$. On the other hand, let U be a non-empty open set. Choose $V \in \bigcup \{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}$ such that $V \setminus M_V \subseteq U$. If $Y \in d^0(\mathcal{V}^*)$, then there exists $W \subseteq V$ such that $W \cap Y = \emptyset$ and $W \in \bigcup \{\mathcal{P}_\alpha : \alpha < \kappa(\prod X_i)\}$. Since $W \setminus M_W \subseteq V \setminus M_V \subseteq U$ we conclude that $Y \cap W \setminus M_W = \emptyset$. $\square$

One can find the next theorem in the paper by W. Kulpa and A. Szymański [8]. It is presented with a proof in [1].

Theorem. *Let $\mathcal{W}$ be a collection of families consisting of open subsets of a topological space Y . Suppose that: $\bigcup \mathcal{W}$ is a π -base; any family in $\mathcal{W}$ consists of pairwise disjoint sets; $|\mathcal{W}| < \tau$, where τ is a regular cardinal number; each set belonging to $\bigcup \mathcal{W}$ contains τ many pairwise disjoint open sets. Then there exists an increasing family of nowhere dense subsets $\{Y_\alpha : \alpha < \tau\}$ such that $\bigcup \{Y_\alpha : \alpha < \tau\} = Y$.*

Thus, we can estimate $\text{cov}(d^0(\mathcal{V}))$, more accurate than these in [6].

Theorem 18. *If sets X_i are countable, then*

$$\text{add}(d^0(\mathcal{V}^*)) \leq \text{cov}(d^0(\mathcal{V}^*)) \leq \text{add}(d^0(\mathcal{V}^*))^+.$$

Proof. Obviously, if $\kappa(\prod X_i) = \mathfrak{c}$, then $\mathfrak{c} = \text{add}(d^0(\mathcal{V}^*)) = \text{cov}(d^0(\mathcal{V}^*))$. Suppose that $\kappa(\prod X_i) < \mathfrak{c}$, then the above theorem, i.e. the theorem by Kulpy and Szymański, works. Let $\mathcal{W}$ be a base matrix. Put $\tau = \kappa(\prod X_i)^+$. Then $\bigcup \mathcal{W}$ is a π -base for the topology generated by itself on $\prod X_i$. Each $V \in \bigcup \mathcal{W}$ contains $\mathfrak{c}$ -many elements of $\bigcup \mathcal{W}$. By Corollary 9 and Lemma 17 and the theorem by Kulpy and Szymański we obtain $\text{cov}(d^0(\mathcal{V}^*)) \leq \kappa(\prod X_i)^+$. We are done, since Theorem 15. $\square$

Thus, if sets X_i are countable, then $\text{cov}(d^0(\mathcal{V})) \leq \text{cov}(d^0(\mathcal{V}^*))$. If sets X_i are finite, then

$$\text{add}(d^0(\mathcal{V})) \leq \text{cov}(d^0(\mathcal{V})) \leq \text{add}(d^0(\mathcal{V}))^+.$$

REFERENCES

- [1] B. Balcar, J. Pelant and P. Simon, *The space of ultrafilters on N covered by nowhere dense sets*, Fund. Math. 110 (1980), no. 1, 11 - 24.
- [2] B. Balcar and P. Simon, *Disjoint refinement*, Handbook of Boolean algebras, North-Holland, Amsterdam, (1989) Vol. 2, 333 - 388.
- [3] J. Brendle, *Strolling through paradise*, Fund. Math. 148 (1995), no. 1, 1 - 25.
- [4] J. Hadamard, *Sur les caracteres de convergence des series a termes positifs et sur les fonctions indefiniment croissantes*, Acta Mathematica 18 (1894), 319 - 336.
- [5] T. Jech, *Set theory*, The third millennium edition, revised and expanded. Springer Monographs in Mathematics. Springer-Verlag, Berlin, (2003).
- [6] P. Kalembe and Sz. Plewik and A. Wojciechowska, *On the ideal (v^0)* , Cent. Eur. J. Math. 6 (2008), no. 2, 218 - 227.
- [7] A. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics 156, Springer-Verlag, New York, (1995).
- [8] W. Kulpa and A. Szymański, *Decomposition into nowhere dense sets*, Bull. Acad. Polon. Sci. 25 (1977), 37 - 39.
- [9] M. Kysiak, A. Nowik and T. Weiss, *Special subsets of the reals and tree forcing notions*, Proc. Amer. Math. Soc. 135 (2007), no. 9, 2975 - 2982.
- [10] G. Moran and D. Strauss, *Countable partitions of product spaces*, Mathematika 27 (1980), no. 2, 213 - 224.
- [11] M. Scheepers, *Gaps in $({}^\omega\omega, <)$* , Set theory of the reals (Ramat Gan, 1991), 439 - 561, Israel Math. Conf. Proc., 6, Bar-Ilan Univ., Ramat Gan, (1993).

PIOTR KALEMBA, INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, UL.
BANKOWA 14, 40-007 KATOWICE

E-mail address: `pkalemba@math.us.edu.pl`

SZYMON PLEWIK, INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, UL.
BANKOWA 14, 40-007 KATOWICE

E-mail address: `plewik@math.us.edu.pl`